

Bavencio® (avelumab) (Intravenous)

-E-

Document Number: OHSU HEALTHSERVICES-0417

Last Review Date: 05/02/2024

Date of Origin: 01/07/2019

Dates Reviewed: 01/2019, 04/2019, 07/2019, 10/2019, 01/2020, 04/2020, 07/2020, 09/2020, 01/2021, 04/2021, 07/2021, 10/2021, 01/2022, 05/2022, 07/2022, 10/2022, 01/2023, 05/2023, 07/2023, 10/2023, 01/2024, 05/2024

I. Length of Authorization

Coverage will be provided for 6 months and may be renewed.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Bavencio 200 mg/10mL single-dose vial: 4 vials per 14 days

B. Max Units (per dose and over time) [HCPCS Unit]:

- 80 billable units (800 mg) every 14 days (all indications)

III. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

- Patient is at least 18 years of age, unless otherwise indicated; **AND**

Universal Criteria

- Patient has not received previous therapy with a programmed death (PD-1/PD-L1)-directed therapy (e.g., nivolumab, pembrolizumab, dostarlimab, atezolizumab, durvalumab, cemiplimab, nivolumab/relatlimab, retifanlimab, toripalimab, etc.), unless otherwise specified ^A; **AND**

Merkel Cell Carcinoma (MCC) † ‡ ☐ ^{1,2,4,5,1e,2e}

- Patient is at least 12 years of age; **AND**
- Used as single-agent therapy; **AND**
- Patient has distant metastatic disease

Urothelial Carcinoma (Bladder Cancer) † ‡ 1,4,6,8,16

- Used as single-agent therapy; **AND**
 - Patient has one of the following diagnoses:
 - Locally advanced or metastatic urothelial carcinoma †
 - Muscle invasive bladder cancer with local recurrence or persistent disease in a preserved bladder treated with curative intent ‡
 - Metastatic or local bladder cancer recurrence post cystectomy treated with curative intent ‡
 - Metastatic upper genitourinary (GU) tract tumors ‡
 - Metastatic urothelial carcinoma of the prostate ‡
 - Recurrent or metastatic primary carcinoma of the urethra (*excluding recurrence of stage T3-4 disease or palpable inguinal lymph nodes*) ‡; **AND**
 - Used for disease that progressed during or following platinum-containing chemotherapy*; **OR**
 - Used as first-line maintenance treatment; **AND**
 - Patient has locally advanced or metastatic urothelial carcinoma (inclusive of bladder, upper GU tract, urethra, and/or prostate cancer); **AND**
 - Patient has not progressed with first-line platinum-containing chemotherapy

*** Note:** 6,17,20

- *If patient was progression-free for > 12 months after platinum therapy, consider re-treatment with platinum-based therapy if the patient is still platinum eligible (see below for cisplatin- or platinum-ineligible comorbidities).*
 - *Cisplatin-ineligible comorbidities may include the following: CrCl < 60 mL/min, ECOG PS ≥ 2 or KPS < 70% hearing loss of ≥ 25 decibels (dB) at two contiguous frequencies, or grade ≥ 2 peripheral neuropathy, or NYHA Heart Failure class ≥ 3. Carboplatin may be substituted for cisplatin particularly in those patients with a CrCl < 60 mL/min or an ECOG PS of 2.*
 - *Platinum-ineligible comorbidities may include the following: CrCl < 30 mL/min, ECOG PS ≥ 3, grade ≥ 2 peripheral neuropathy, or NYHA Heart Failure class > 3, etc.*

Renal Cell Carcinoma (RCC) † ‡ 1,4,9,14

- Used in combination with axitinib; **AND**
- Used as first-line therapy; **AND**
- Used for the treatment of advanced, relapsed, or stage IV disease and clear cell histology

Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Φ Orphan Drug

IV. Renewal Criteria ^{Δ 1}

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria identified in section III; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: severe or life-threatening infusion-related reactions, hepatotoxicity, severe immune-mediated adverse reactions (e.g., pneumonitis, hepatitis, colitis, endocrinopathies, nephritis with renal dysfunction, dermatitis/dermatologic adverse reactions, etc.), major adverse cardiovascular events (MACE) when used in combination with axitinib, complications of allogeneic hematopoietic stem cell transplantation (HSCT), etc.
-

^Δ Notes:

- Patients whose tumors, upon re-biopsy, demonstrate a change in actionable mutation (e.g., MSS initial biopsy; MSI-H subsequent biopsy) may be eligible to re-initiate PD-directed therapy and will be evaluated on a case-by-case basis.

V. Dosage/Administration ^{1,13,18}

Indication	Dose
All indications	Administer 800 mg intravenously every 14 days, until disease progression or unacceptable toxicity
<p><u>Dosing should be calculated using actual body weight and not flat dosing (as applicable) based on the following:</u></p> <p>Weight is ≤ 66 kg:</p> <ul style="list-style-type: none"> • Use 600 mg (10mg/kg) IV every 2 weeks <p><i>Note: This information is not meant to replace clinical decision making when initiating or modifying medication therapy and should only be used as a guide. Patient-specific variables should be taken into account.</i></p>	

VI. Billing Code/Availability Information

HCPCS Code:

- J9023 – Injection, avelumab, 10 mg; 1 billable unit = 10 mg

NDC:

- Bavencio 200 mg/10 mL single-dose vial: 44087-3535-xx

VII. References (STANDARD)

1. Bavencio [package insert]. Rockland, MA; EMD Serono, Inc; March 2024. Accessed March 2024.
2. Kaufman HL, Russell J, Hamid O, et al. Avelumab in patients with chemotherapy-refractory metastatic Merkel cell carcinoma: a multicentre, single-group, open-label, phase 2 trial. *Lancet Oncol.* 2016 Oct;17(10):1374-1385.
3. Novakovic AM, Wilkins JJ, Dai H, et al. Changing body weight-based dosing to a flat dose for avelumab in metastatic Merkel cell and advanced urothelial carcinoma. *Clin Pharmacol Ther.* 2019 Sep 25.
4. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) avelumab. National Comprehensive Cancer Network, 2024. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2024.
5. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Merkel Cell Carcinoma, Version 1.2024. National Comprehensive Cancer Network, 2024. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2024.
6. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Bladder Cancer, Version 1.2024. National Comprehensive Cancer Network, 2024. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2024.
7. Gupta S, Sonpavde G, Grivas P, et al. Defining “platinum-ineligible” patients with metastatic urothelial cancer (mUC). *J Clin Oncol.* 2019 Mar 1;37(7_suppl):451.
8. Patel MR, Ellerton J, Infante JR, et al. Avelumab in metastatic urothelial carcinoma after platinum failure (JAVELIN Solid Tumor): pooled results from two expansion cohorts of an open-

- label, phase 1 trial. *Lancet Oncol.* 2018 Jan;19(1):51-64. doi: 10.1016/S1470-2045(17)30900-2. Epub 2017 Dec 5.
9. Motzer RJ, Penkov K, Haanen J, et al. Avelumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma. *N Engl J Med.* 2019 Mar 21;380(12):1103-1115. doi: 10.1056/NEJMoa1816047. Epub 2019 Feb 16.
 10. Fahrenbruch R, Kintzel P, Bott AM, et al. Dose Rounding of Biologic and Cytotoxic Anticancer Agents: A Position Statement of the Hematology/Oncology Pharmacy Association. *J Oncol Pract.* 2018 Mar;14(3):e130-e136.
 11. Hematology/Oncology Pharmacy Association (2022). Intravenous Cancer Drug Waste Issue Brief. Retrieved from: https://www.hoparx.org/documents/65/HOPA_Drug_Waste_Issue_Brief_-_Updated_01.19.22_FINAL.pdf
 12. Bach PB, Conti RM, Muller RJ, et al. Overspending driven by oversized single dose vials of cancer drugs. *BMJ.* 2016 Feb 29;352:i788.
 13. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Gestational Trophoblastic Neoplasia. Version 1.2024. National Comprehensive Cancer Network, 2024. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2024.
 14. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Kidney Cancer. Version 1.2024. National Comprehensive Cancer Network, 2024. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2024.
 15. You B, Bolze PA, Lotz JP, et al. Avelumab in Patients With Gestational Trophoblastic Tumors With Resistance to Single-Agent Chemotherapy: Cohort A of the TROPHIMMUN Phase II Trial. *J Clin Oncol.* 2020 Sep 20;38(27):3129-3137. doi: 10.1200/JCO.20.00803. Epub 2020 Jul 27.
 16. Powles T, Park SH, Voog E, et al. Avelumab Maintenance Therapy for Advanced or Metastatic Urothelial Carcinoma. *N Engl J Med.* 2020 Sep 24;383(13):1218-1230. doi: 10.1056/NEJMoa2002788. Epub 2020 Sep 18. PMID: 32945632.
 17. Bellmunt, J. (2024). Treatment of metastatic urothelial cancer of the bladder and urinary tract. In Lerner SP, Shah S (Eds.), *UptoDate*. Last updated February 7, 2024. Accessed February 12, 2024. Available from https://www.uptodate.com/contents/treatment-of-metastatic-urothelial-cancer-of-the-bladder-and-urinary-tract?search=cisplatin%20ineligible&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1.

18. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Uterine Neoplasms. Version 1.2024. National Comprehensive Cancer Network, 2024. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2024.
19. Galsky MD, Balar AV, Black PC, et al. Society for Immunotherapy of Cancer (SITC) clinical practice guideline on immunotherapy for the treatment of urothelial cancer. *Journal for ImmunoTherapy of Cancer* 2021;9:e002552. doi: 10.1136/jitc-2021-002552.
20. Gupta S, Bellmunt J, Plimack ER, et al. Defining “platinum-ineligible” patients with metastatic urothelial cancer (mUC). *J Clin Oncol*. 2022 June 1;40(16_suppl):4577.
21. You B, Bolze PA, Lotz JP, et al. Avelumab in patients with gestational trophoblastic tumors with resistance to polychemotherapy: Cohort B of the TROPHIMMUN phase 2 trial. *Gynecol Oncol*. 2023 Jan;168:62-67. doi: 10.1016/j.ygyno.2022.11.005.

VIII. References (ENHANCED)

1. D'Angelo SP, Russell J, Lebbé C, et al. Efficacy and Safety of First-line Avelumab Treatment in Patients With Stage IV Metastatic Merkel Cell Carcinoma: A Preplanned Interim Analysis of a Clinical Trial. *JAMA Oncol*. 2018;4(9):e180077.
2. Kaufman HL, Russell JS, Hamid O, et al. Updated efficacy of avelumab in patients with previously treated metastatic Merkel cell carcinoma after ≥1 year of follow-up: JAVELIN Merkel 200, a phase 2 clinical trial. *J Immunother Cancer*. 2018;6(1):7. Published 2018 Jan 19. doi:10.1186/s40425-017-0310-x.
3. Nghiem PT, Bhatia S, Lipson EJ, et al. PD-1 Blockade with Pembrolizumab in Advanced Merkel-Cell Carcinoma. *N Engl J Med*. 2016;374(26):2542-52.
4. Topalian SL, Bhatia S, Hollebecque A, et al. Abstract CT074: Non-comparative, open-label, multiple cohort, phase 1/2 study to evaluate nivolumab (NIVO) in patients with virus-associated tumors (CheckMate 358): Efficacy and safety in Merkel cell carcinoma (MCC). *Cancer Res* 2017;77(13 Suppl):Abstract nr CT074.
5. Bellmunt J, de Wit R, Vaughn DJ, et al. Pembrolizumab as Second-Line Therapy for Advanced Urothelial Carcinoma. *N Engl J Med*. 2017;376(11):1015-1026.
6. Apolo AB, Infante JR, Balmanoukian A, et al. Avelumab, an Anti-Programmed Death-Ligand 1 Antibody, In Patients With Refractory Metastatic Urothelial Carcinoma: Results From a Multicenter, Phase Ib Study. *J Clin Oncol*. 2017;35(19):2117-2124.
7. Sharma P, Retz M, Siefker-Radtke A, et al. Nivolumab in metastatic urothelial carcinoma after platinum therapy (CheckMate 275): a multicentre, single-arm, phase 2 trial. *Lancet Oncol*. 2017 Mar;18(3):312-322.

8. Rini BI, Plimack ER, Stus V, et al. Pembrolizumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma. *N Engl J Med*. 2019 Mar 21;380(12):1116-1127.
9. Motzer RJ, Tannir NM, McDermott DF, et al. Nivolumab plus Ipilimumab versus Sunitinib in Advanced Renal-Cell Carcinoma. *N Engl J Med*. 2018;378(14):1277–1290.
10. Siefker-Radtke AO, Necchi A, Park SH, et al. First results from the primary analysis population of the phase 2 study of erdafitinib (ERDA; JNJ-42756493) in patients (pts) with metastatic or unresectable urothelial carcinoma (mUC) and FGFR alterations (FGFRalt). *J Clin Oncol* 2018;36(15_suppl):4503.
11. Powles T, Park SH, Voog E, et al. Maintenance avelumab + best supportive care (BSC) versus BSC alone after platinum-based first-line (1L) chemotherapy in advanced urothelial carcinoma (UC): JAVELIN Bladder 100 phase III interim analysis. *J Clin Oncol*. 2020;38(18_suppl):LBA1-LBA1.
12. You B, Bolze PA, Lotz JP, et al. Avelumab in Patients With Gestational Trophoblastic Tumors With Resistance to Single-Agent Chemotherapy: Cohort A of the TROPHIMMUN Phase II Trial. *J Clin Oncol*. 2020 Sep 20;38(27):3129-3137. doi: 10.1200/JCO.20.00803. Epub 2020 Jul 27.
13. Ghorani E, Kaur B, Fisher RA, et al. Pembrolizumab is effective for drug-resistant gestational trophoblastic neoplasia. *Lancet*. 2017 Nov 25;390(10110):2343-2345. doi: 10.1016/S0140-6736(17)32894-5.
14. Choueiri TK, Powles T, Burotto M, et al. Nivolumab plus Cabozantinib versus Sunitinib for Advanced Renal-Cell Carcinoma. *N Engl J Med*. 2021 Mar 4;384(9):829-841. doi: 10.1056/NEJMoa2026982.
15. Motzer R, Alekseev B, Rha SY, et al. Lenvatinib plus Pembrolizumab or Everolimus for Advanced Renal Cell Carcinoma. *N Engl J Med*. 2021 Apr 8;384(14):1289-1300. doi: 10.1056/NEJMoa2035716. Epub 2021 Feb 13.
16. Choueiri TK, Hessel C, Halabi S, et al. Cabozantinib versus sunitinib as initial therapy for metastatic renal cell carcinoma of intermediate or poor risk (Alliance A031203 CABOSUN randomised trial): Progression-free survival by independent review and overall survival update [published correction appears in *Eur J Cancer*. 2018 Nov;103:287]. *Eur J Cancer*. 2018;94:115-125. doi:10.1016/j.ejca.2018.02.012.
17. Konstantinopoulos PA, Luo W, Liu JF, et al. Phase II Study of Avelumab in Patients With Mismatch Repair Deficient and Mismatch Repair Proficient Recurrent/Persistent Endometrial Cancer. *J Clin Oncol*. 2019 Oct 20;37(30):2786-2794. doi: 10.1200/JCO.19.01021. Epub 2019 Aug 28. PMID: 31461377.
18. You B, Bolze P, Lotz J, et al. 273 Avelumab in patients with gestational trophoblastic tumors resistant to polychemotherapy: efficacy outcomes of cohort B of TROPHIMMUN phase II trial. *International Journal of Gynecologic Cancer* 2021;31:A344.
19. Powles T, Rosenberg JE, Sonpavde GP, et al. Enfortumab Vedotin in Previously Treated Advanced Urothelial Carcinoma. *N Engl J Med*. 2021 Mar 25;384(12):1125-1135. doi: 10.1056/NEJMoa2035807. Epub 2021 Feb 12.

20. Magellan Rx Management. Bavencio Clinical Literature Review Analysis. Last updated March 2024. Accessed March 2024.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C4A.0	Merkel cell carcinoma of lip
C4A.10	Merkel cell carcinoma of eyelid, including canthus
C4A.111	Merkel cell carcinoma of right upper eyelid, including canthus
C4A.112	Merkel cell carcinoma of right lower eyelid, including canthus
C4A.121	Merkel cell carcinoma of left upper eyelid, including canthus
C4A.122	Merkel cell carcinoma of left lower eyelid, including canthus
C4A.20	Merkel cell carcinoma of unspecified ear and external auricular canal
C4A.21	Merkel cell carcinoma of right ear and external auricular canal
C4A.22	Merkel cell carcinoma of left ear and external auricular canal
C4A.30	Merkel cell carcinoma of unspecified part of face
C4A.31	Merkel cell carcinoma of nose
C4A.39	Merkel cell carcinoma of other parts of face
C4A.4	Merkel cell carcinoma of scalp and neck
C4A.51	Merkel cell carcinoma of anal skin
C4A.52	Merkel cell carcinoma of skin of breast
C4A.59	Merkel cell carcinoma of other part of trunk
C4A.60	Merkel cell carcinoma of unspecified upper limb, including shoulder
C4A.61	Merkel cell carcinoma of right upper limb, including shoulder
C4A.62	Merkel cell carcinoma of left upper limb, including shoulder
C4A.70	Merkel cell carcinoma of unspecified lower limb, including hip
C4A.71	Merkel cell carcinoma of right lower limb, including hip
C4A.72	Merkel cell carcinoma of left lower limb, including hip
C4A.8	Merkel cell carcinoma of overlapping sites
C4A.9	Merkel cell carcinoma, unspecified
C61	Malignant neoplasm of prostate
C64.1	Malignant neoplasm of right kidney, except renal pelvis
C64.2	Malignant neoplasm of left kidney, except renal pelvis
C64.9	Malignant neoplasm of unspecified kidney, except renal pelvis
C65.1	Malignant neoplasm of right renal pelvis
C65.2	Malignant neoplasm of left renal pelvis
C65.9	Malignant neoplasm of unspecified renal pelvis
C66.1	Malignant neoplasm of right ureter

ICD-10	ICD-10 Description
C66.2	Malignant neoplasm of left ureter
C66.9	Malignant neoplasm of unspecified ureter
C67.0	Malignant neoplasm of trigone of bladder
C67.1	Malignant neoplasm of dome of bladder
C67.2	Malignant neoplasm of lateral wall of bladder
C67.3	Malignant neoplasm of anterior wall of bladder
C67.4	Malignant neoplasm of posterior wall of bladder
C67.5	Malignant neoplasm of bladder neck
C67.6	Malignant neoplasm of ureteric orifice
C67.7	Malignant neoplasm of urachus
C67.8	Malignant neoplasm of overlapping sites of bladder
C67.9	Malignant neoplasm of bladder, unspecified
C68.0	Malignant neoplasm of urethra
C7B.1	Secondary Merkel cell carcinoma
D09.0	Carcinoma in situ of bladder
Z85.51	Personal history of malignant neoplasm of bladder
Z85.59	Personal history of malignant neoplasm of other urinary tract organ
Z85.821	Personal history of Merkel cell carcinoma

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC