OHSU Health	Page	S Provider's Orders	ACCOUNT NO. MED. REC. NO. NAME BIRTHDATE Patient Identification IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.			
Allergies:	kg	Height:				
Treatment Start Date: Patient to follow up with provider on date:						

GUIDELINES FOR ORDERING

1. Send FACE SHEET and H&P or most recent chart note.

2. Provider must order and obtain a ferritin prior to patient being scheduled for iron infusion. Labs drawn date: _____

NURSING ORDERS:

- 1. TREATMENT PARAMETER Hold treatment and notify provider if Ferritin greater than 300 ng/mL.
- 2. Instruct patient to set follow up appointment with provider for follow up labs.
- 3. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

MEDICATIONS:

iron sucrose (VENOFER): (must check one)

- 200 mg in sodium chloride 0.9% 100 mL, intravenous, ONCE, over 30 minutes, x 5 doses over 14 days
- 300 mg in sodium chloride 0.9% 250 mL, intravenous, ONCE, over 1.5 hours, x 3 doses (administered every 2 to 3 days)

AS NEEDED MEDICATIONS:

1. sodium chloride 0.9%, 500 mL, intravenous, AS NEEDED x1 dose for vein discomfort. Give concurrently with iron sucrose

HYPERSENSITIVITY MEDICATIONS:

- 1. NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
- 2. NURSING COMMUNICATION Avoid intravenous or oral diphenhydrAMINE, move to next option in the algorithm. Adverse effects of diphenhydrAMINE may overlap with IV iron adverse effects such as flushing, hypotension, tachycardia.
- 3. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

ONLINE 11/2024 [supersedes 03/2024]

ONLINE 11/2024 [supersedes 03/2024]	

Tage 2 01 2	
	Patient Identification
ALL ORDERS MUST BE MARKED	IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.
By signing below, I represent the following:	
I am responsible for the care of the patient (who is	s identified at the top of this form);
I hold an active, unrestricted license to practice me	
	re to patient and where you are currently licensed. Specify
state if not Oregon);	
My physician license Number is #	(MUST BE COMPLETED TO BE A VALID
	e of practice and authorized by law to order Infusion of the
medication described above for the patient identifi	1 2

Provider signature:	Date/Time:	
Printed Name:	Phone:	Fax:

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

Beaverton
 OHSU Knight Cancer Institute
 15700 SW Greystone Court
 Beaverton, OR 97006
 Phone number: 971-262-9000
 Fax number: 503-346-8058

□ Gresham

Legacy Mount Hood campus Medical Office Building 3, Suite 140 24988 SE Stark Gresham, OR 97030 Phone number: 971-262-9500 Fax number: 503-346-8058

□ NW Portland

Legacy Good Samaritan campus Medical Office Building 3, Suite 150 1130 NW 22nd Ave Portland, OR 97210 Phone number: 971-262-9600 Fax number: 503-346-8058

□ Tualatin

Legacy Meridian Park campus Medical Office Building 2, Suite 140 19260 SW 65th Ave Tualatin, OR 97062 Phone number: 971-262-9700 Fax number: 503-346-8058

PO-8066

Infusion orders located at: www.ohsuknight.com/infusionorders

Oregon Health & Science University Hospital and Clinics Provider's Orders ACCOUNT NO. OHSU Health ADULT AMBULATORY INFUSION ORDER Iron Sucrose (VENOFER) Infusion MED. REC. NO. Page 2 of 2 BIRTHDATE