Oregon Health & Science University Hospital and Clinics Provider's Orders	ACCOUNT NO. MED. REC. NO. NAME BIRTHDATE			
Pentamidine (PENTAM) Infusion				
Page 1 of 3	Patient Identification			
Patient Identification ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.				
Weight:kg Height:	cm			
Allergies:				
Diagnosis Code:				
Treatment Start Date: Patient to follow up with provider on date:				
This plan will expire after 365 days at which	time a new order will need to be placed			
 GUIDELINES FOR ORDERING Send FACE SHEET and H&P or most recent chart note. 12 Lead ECG should be completed prior to treatment with pentamidine. Results MUST be faxed with this order set to be kept on record within the infusion pharmacy's electronic medical record. Avoid use in patients with diagnosed or suspected congenital long QT syndrome. Use with caution in patients with pre-existing hypotension. Severe hypotension including fatalities, has been observed even after a single dose. Use with caution in patients with pre-existing cardiovascular disease, diabetes mellitus, or hypocalcemia. Use with caution in patients receiving nephrotoxic drugs such as aminoglycosides, amphotericin B, cisplatin, foscarnet, or vancomycin. 				
OTHER:				
12 Lead ECG, routine, ONCE every	weeks			
LABS: CMP (includes blood glucose), Routine, ONCE, every (visit)(days)(weeks)(months) – Circle One Glucose (serum), Routine, ONCE, every (visit)(days)(weeks)(months) – Circle One CBC with differential, Routine, ONCE, weekly during therapy CBC with differential, Routine, ONCE, every (visit)(days)(weeks)(months) – Circle One Labs already drawn. Date:				
MEDICATIONS:				
pentamidine (PENTAM) in dextrose 5% 25	0 mL, intravenous, ONCE			

- $\Box 3 mg/kg = \underline{\qquad} mg$ $\Box 4 mg/kg = \underline{\qquad} mg$

Infuse slowly over 1-2 hours. Vesicant. Flush line with D5W before and after infusion.

Interval: (must check one)

- □ Once
- \Box Once daily x _____ doses
- □ ____ times per week x ____ doses □ Monthly x _____ doses

ONLINE 11/2024 [supersedes 02/2023]

OHSU ADULT AMBULATORY INFUSION ORDER Health Pentamidine (PENTAM) Infusion	ACCOUNT NO. MED. REC. NO. NAME BIRTHDATE
Page 2 of 3	
	Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

NURSING ORDERS:

- 1. Review patient's SCr, BUN, calcium, and blood glucose during each visit. Notify provider if laboratory values are abnormal.
- 2. VITAL SIGNS Monitor patient's blood pressure for hypotension during and after infusion
- 3. Instruct patient to lie supine during the infusion. Patient should rise slowly after administration to avoid dizziness and other potentially severe hypotensive effects.
- 4. This medication is a vesicant. Avoid extravasation. Assess catheter position before and during infusion
- 5. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

AS NEEDED MEDICATIONS:

Г

1. prochlorperazine (COMPAZINE) tablet, 10mg, oral, AS NEEDED, x1 doses for nausea/vomiting

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*); I hold an active, unrestricted license to practice medicine in: *Oregon* (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is #______ (MUST BE COMPLETED TO BE A VALID

PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature:		Date/Time:	
Printed Name:	Phone:		Fax:

Oregon Health & Science University Hospital and Clinics Provider's Orders	ACCOUNT NO.
OHSU ADULT AMBULATORY INFUSION ORDER	MED. REC. NO.
Health Pentamidine (PENTAM) Infusion	NAME
Page 3 of 3	BIRTHDATE
	Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (\checkmark) TO BE ACTIVE.

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

□ Beaverton

OHSU Knight Cancer Institute 15700 SW Greystone Court Beaverton, OR 97006 Phone number: 971-262-9000 Fax number: 503-346-8058

□ Gresham

Legacy Mount Hood campus Medical Office Building 3, Suite 140 24988 SE Stark Gresham, OR 97030 Phone number: 971-262-9500 Fax number: 503-346-8058

□ NW Portland

Legacy Good Samaritan campus Medical Office Building 3, Suite 150 1130 NW 22nd Ave. Portland, OR 97210 Phone number: 971-262-9600 Fax number: 503-346-8058

□ Tualatin

Legacy Meridian Park campus Medical Office Building 2, Suite 140 19260 SW 65th Ave. Tualatin, OR 97062 Phone number: 971-262-9700 Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders