



Oregon Health & Science University
Hospital and Clinics Provider's Orders

PO7071



ADULT AMBULATORY INFUSION ORDER
Antibiotic Therapy
(Cephalosporin, Fluoroquinolone, and Others)

Page 1 of 4

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET** and **H&P** or most recent chart note.

LABS:

- CBC with differential, Routine, ONCE, every _____ (visit)(days)(weeks)(months) – Circle One
- CMP, Routine, ONCE, every _____ (visit)(days)(weeks)(months) – Circle One
- Labs already drawn. Date: _____

NURSING ORDERS:

1. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declothing (alteplase), and/or dressing changes
2. In the case of sulfamethoxazole/trimethoprim (BACTRIM), flush IV line with 5 mL dextrose 5% before and after each infusion.

MEDICATIONS:

Cephalosporins:

- ceFAZolin 500 mg in sodium chloride 0.9% 100 mL IV, ONCE over 20-40 minutes
- ceFAZolin 1 gram in sodium chloride 0.9% 100 mL IV, ONCE over 20-40 minutes
- ceFAZolin 6 grams over 1 day in sodium chloride 0.9% 151.2 mL IV, ONCE over 24 hours, continuous infusion via CADD (OHSU only)

- ceFEPime 1 gram in sodium chloride 0.9% 50 mL IV, ONCE over 30 minutes
- ceFEPime 2 grams in sodium chloride 0.9% 50 mL IV, ONCE over 30 minutes
- ceFEPime 4 grams over 1 day in sodium chloride 0.9% 100.8 mL IV, ONCE over 24 hours, continuous infusion via CADD (OHSU only)
- ceFEPime 6 grams over 1 day in sodium chloride 0.9% 151.2 mL IV, ONCE over 24 hours, continuous infusion via CADD (OHSU only)

- ceftAZidime 1 gram in sodium chloride 0.9% 100 mL IV, ONCE over 15-30 minutes
- ceftAZidime 2 grams in sodium chloride 0.9% 100 mL IV, ONCE over 15-30 minutes
- ceftAZidime 6 grams over 1 day in sodium chloride 0.9% 151.2 mL IV, ONCE over 24 hours, continuous infusion via CADD (OHSU only)



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
Antibiotic Therapy
(Cephalosporin, Fluoroquinolone, and Others)

Page 2 of 4

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

- cefTRIAxone 1 gram
- cefTRIAxone 2 grams

In sodium chloride 0.9% 50 mL IV over 30 minutes, or 100 mg/mL IV push over 5 minutes per infusion facility practice.

Interval: (must check one)

- ONCE
- Daily x ____ doses

Fluoroquinolones:

- ciprofloxacin 200 mg in sodium chloride 0.9% 200 mL IV, ONCE over 60 minutes
- ciprofloxacin 400 mg in sodium chloride 0.9% 200 mL IV, ONCE over 60 minutes

- levoFLOxacin 250 mg in sodium chloride 0.9% 50 mL IV, ONCE over 60 minutes
- levoFLOxacin 500 mg in sodium chloride 0.9% 100 mL IV, ONCE over 60 minutes
- levoFLOxacin 750 mg in sodium chloride 0.9% 150 mL IV, ONCE over 90 minutes

Interval: (must check one)

- ONCE
- Daily x ____ doses

Other:

- azithromycin 250 mg in sodium chloride 0.9% 250 mL IV, ONCE over 60 minutes
- azithromycin 500 mg in sodium chloride 0.9% 250 mL IV, ONCE over 60 minutes

- clindamycin 600 mg in sodium chloride 0.9% 50 mL IV, ONCE over 30 minutes
- clindamycin 900 mg in sodium chloride 0.9% 50 mL IV, ONCE over 30 minutes

- doxycycline 100 mg in sodium chloride 0.9% 250 mL IV, ONCE over 60 minutes
- doxycycline 200 mg in sodium chloride 0.9% 250 mL IV, ONCE over 60 minutes

- sulfamethoxazole/trimethoprim 5 mg/kg = _____ mg in **dextrose 5%** IV, ONCE over 60-90 minutes

- Other (drug, dose, route):** _____
(Pharmacist to confirm availability)

Interval: (must check one)

- ONCE
- Daily x ____ doses



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
Antibiotic Therapy
(Cephalosporin, Fluoroquinolone, and Others)

Page 3 of 4

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

FOR InfuSystem™ AMBULATORY PUMP USE (OHSU only; hook up at infusion location):

Duration:

_____ days

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____



**Oregon Health & Science University
Hospital and Clinics Provider's Orders**

ADULT AMBULATORY INFUSION ORDER
Antibiotic Therapy
(Cephalosporin, Fluoroquinolone, and Others)

Page 4 of 4

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Central Intake:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:

Beaverton

OHSU Knight Cancer Institute
15700 SW Greystone Court
Beaverton, OR 97006

Phone number: 971-262-9000

Fax number: 503-346-8058

NW Portland

Legacy Good Samaritan campus
Medical Office Building 3, Suite 150
1130 NW 22nd Ave
Portland, OR 97210

Phone number: 971-262-9600

Fax number: 503-346-8058

Gresham

Legacy Mount Hood campus
Medical Office Building 3, Suite 140
24988 SE Stark
Gresham, OR 97030

Phone number: 971-262-9500

Fax number: 503-346-8058

Tualatin

Legacy Meridian Park campus
Medical Office Building 2, Suite 140
19260 SW 65th Ave
Tualatin, OR 97062

Phone number: 971-262-9700

Fax number: 503-346-8058

Infusion orders located at: www.ohsuknight.com/infusionorders