

# Oregon Health & Science University Hospital and Clinics Provider's Orders



ADULT AMBULATORY INFUSION ORDER remdesivir (VEKLURY) infusion Infusion
Page 1 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

# ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.

Weight:kg	Height: _	cm	
Allergies:			
Diagnosis Code:			
Treatment Start Date:		Patient to follow up with provider on date:	

#### **GUIDELINES FOR ORDERING**

- 1. Send FACE SHEET and H&P or most recent chart note.
- 2. COVID-19 therapies are NOT authorized in patients with known hypersensitivity to any ingredient of the treatment.
- 3. To be eligible for treatment, patient must meet the criteria below:
  - a. At least 3 kg or greater
  - b. Documented positive PCR or antigen test
  - c. Symptomatic COVID-19 infections
  - d. Within 7 days of COVID-19 symptom onset
  - e. Not on supplemental oxygen or requiring more oxygen than baseline if on chronic oxygen

## AND at least one of the following

- □ Body mass index is 25 or greater
- □ Pregnancy
- ☐ Chronic kidney disease
- □ Diabetes
- ☐ Immunosuppressive disease or Immunosuppressive treatment
- □ Sickle cell disease
- ☐ Cardiovascular disease (including congenital heart disease) or hypertension
- ☐ Chronic lung diseases
- □ Neurodevelopmental disorders or medical-related technological dependence
- ☐ Age greater than or equal to 65 years
- ☐ Cardiovascular disease, or hypertension, or COPD/other Chronic respiratory disease
- □ 12-17 years of age AND BMI greater than 85th percentile for age and gender based on CDC growth chart, OR sickle cell disease, OR congenital/acquired heart disease, OR medical-related technological dependence, OR neurodevelopmental disorder, OR medical related technological dependence, OR asthma/reactive airway/other chronic respiratory disease requiring daily medication for control
- 4. Hepatitis B (Hep B surface antigen and core antibody total) screening must be completed prior to initiation of treatment and the patient should not be infected. Please send results with order.
- 5. A Tuberculin test must have been placed and read as negative prior to initiation of treatment (PPD or QuantiFERON Gold blood test). Please send results with order. If result is indeterminate, a follow up chest X-ray must be performed to rule out TB. Please send results with order.

<sup>\*\*</sup>This plan will expire after 365 days at which time a new order will need to be placed\*\*



# Oregon Health & Science University Hospital and Clinics Provider's Orders

OHSU ADULT AMBULATORY INFUSION ORDER Health remdesivir (VEKLURY) infusion

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.

ı	Λ	DC	

Liver Set (AST, ALT, BILI TOTAL, BILI DIRECT, ALK PHOS, ALB, PROT TOTAL), Routine, ONCE,
every (visit)(days)(weeks)(months) - Circle One
Labs already drawn. Date:

#### **NURSING ORDERS:**

- 1. TREATMENT PARAMETERS Hold treatment and notify provider if patient does not meet criteria above in Guidelines for Ordering, or if treatment day #1 ALT is greater than or equal to 10 x ULN.
- 2. Contact provider if any concerns of adverse drug reactions.
- 3. Monitor patient during administration and observe for hypersensitivity reactions, including anaphylaxis, for 1 hour after administration.
- 4. May leave PIV in place for consecutive day infusions if clinically indicated and appropriate.
- 5. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

#### **MEDICATIONS:**

remdesivir (VEKLURY) in sodiu	um chloride 0.9%, intravenous, ONCE, over 30 minutes
☐ 200 mg ONCE on day 1	, followed by 100 mg DAILY on days 2 and 3
□ 100 mg DAILY x	dose(s) to complete previously started treatment course

## HYPERSENSITIVITY MEDICATIONS:

- NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the
  infusion and notify provider immediately. Administer emergency medications per the Treatment
  Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for
  symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 3. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

### **AS NEEDED MEDICATIONS:**

- 1. acetaminophen (TYLENOL) tablet, 650 mg, oral, EVERY 4 HOURS AS NEEDED for fever
- 2. diphenhydrAMINE (BENADRYL) capsule, 25 mg, oral, EVERY 4 HOURS AS NEEDED for itching
- 3. sodium chloride 0.9% solution, intravenous, 500mL, AS NEEDED x1 dose, for TNF-alpha inhibitor infusion tolerability. Give concurrently with TNF-alpha inhibitor



# Oregon Health & Science University Hospital and Clinics Provider's Orders

OHSU ADULT AMBULATORY INFUSION ORDER remdesivir (VEKLURY) infusion

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.

By signing below, I represent the following:  I am responsible for the care of the patient (who is identified at the top of this form);  I hold an active, unrestricted license to practice medicine in:   Oregon   (check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon);						
My physician license Number is #	(MUST BE COM	MPLETED TO BE A VALID				
<u>PRESCRIPTION</u> ); and I am acting within r medication described above for the patient		ed by law to order Infusion of the				
Provider signature:	Date/Time	Date/Time:				
Printed Name:	Phone:	Fax:				
OLC Central Intake Nurse:  Phone: 971-262-9645 (providers only) Fax  Please check the appropriate box for the		tion:				
□ Beaverton OHSU Knight Cancer Institute 15700 SW Greystone Court Beaverton, OR 97006 Phone number: 971-262-9000 Fax number: 503-346-8058	□ <b>NW Portland</b> Legacy Good Sar	maritan campus iilding 3, Suite 150 ve. 10 <mark>71-262-9600</mark>				
☐ Gresham Legacy Mount Hood campus Medical Office Building 3, Suite 140 24988 SE Stark Gresham, OR 97030 Phone number: 971-262-9500 Fax number: 503-346-8058	☐ Tualatin  Legacy Meridian  Medical Office Bu  19260 SW 65th A  Tualatin, OR 9700  Phone number: Solution	nilding 2, Suite 140 Ne. 62 <mark>971-262-9700</mark>				

Infusion orders located at: <a href="https://www.ohsuknight.com/infusionorders">www.ohsuknight.com/infusionorders</a>